arXiv:1601.02864v1 [math.CO] 12 Jan 2016

# TABLES OF SUBSPACE CODES

DANIEL HEINLEIN, MICHAEL KIERMAIER, SASCHA KURZ, AND ALFRED WASSERMANN*

ABSTRACT. The main problem of subspace coding asks for the maximum possible cardinality of a subspace code with minimum distance at least $d$ over $\mathbb{F}_q^n$, where the dimensions of the codewords, which are vector spaces, are contained in $K \subseteq \{0, 1, \ldots, n\}$. In the special case of $K = \{k\}$ one speaks of constant dimension codes. Since this emerging field is very prosperous on the one hand side and there are a lot of connections to classical objects from Galois geometry it is a bit difficult to keep or to obtain an overview about the current state of knowledge. To this end we have implemented an on-line database of the (at least to us) known results at `subspacecodes.uni-bayreuth.de`. The aim of this recurrently updated technical report is to provide a user guide how this technical tool can be used in research projects and to describe the so far implemented theoretic and algorithmic knowledge.

**Keywords:** Galois geometry, subspace codes, partial spreads, constant dimension codes
**MSC:** 51E23; 05B40, 11T71, 94B25

## 1. INTRODUCTION

The seminal paper by Kötter and Kschischang [15] started the interest in subspace codes which are sets of subspaces of the $\mathbb{F}_q$-vector space $\mathbb{F}_q^n$. Two widely used distance measures for subspace codes (motivated by an information-theoretic analysis of the Koetter-Kschischang-Silva model, see e.g. [21]) are the *subspace distance*

$$d_S(U,W) := \dim(U+W) - \dim(U \cap W) = 2 \cdot \dim(U+W) - \dim(U) - \dim(W)$$

and the *injection distance*

$$d_I(U,W) := \max\{\dim(U), \dim(W)\} - \dim(U \cap W),$$

where $U$ and $W$ are subspaces of $\mathbb{F}_q^n$. The two metrics are equivalent, i.e. it is known that $d_I(U,W) \leq d_S(U,W) \leq 2d_I(U,W)$.

The set of all $k$-dimensional subspaces of an $\mathbb{F}_q$-vector space $V$ will be denoted by $\begin{bmatrix} V \\ k \end{bmatrix}_q$. For $n = \dim(V)$, its cardinality is given by the Gaussian binomial coefficient

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \begin{cases} \frac{(q^n-1)(q^{n-1}-1)\cdots(q^{n-k+1}-1)}{(q^k-1)(q^{k-1}-1)\cdots(q-1)} & \text{if } 0 \leq k \leq n; \\ 0 & \text{otherwise.} \end{cases}$$

A set $\mathcal{C}$ of subspaces of $V$ is called a *subspace code*. The *minimum distance* of $\mathcal{C}$ is given by $d = \min\{d_S(U,W) \mid U, W \in \mathcal{C}, U \neq W\}$. If the dimensions of the *codewords*, i.e. the elements of $\mathcal{C}$ are contained in some set $K \subseteq \{1, \ldots, n\}$, $\mathcal{C}$ is called a $(n, \#\mathcal{C}, d; K)_q$ subspace code. In the unrestricted case $K = \{0, \ldots, n\}$ we use the notation $(n, \#\mathcal{C}, d)_q$ subspace code. In the other extreme case $K = \{k\}$, we use the notation $(n, \#\mathcal{C}, d)_q$ and call $\mathcal{C}$ a *constant dimension code*.

For fixed ambient parameters $q$, $K$ and $d$, the *main problem of subspace coding* asks for the determination of the maximum possible size $A_q^S(n, d; K) := M$ of an $(n, M, d)_q$ subspace code and – as a refinement – the classification of all corresponding optimal codes up to isomorphism. Again, the simplified notations $A_q^S(n, d)$ and $A_q^S(n, d; k)$ are used for the unrestricted case $K = \{0, \ldots, n\}$ and the constant

* The work of the authors was supported by the ICT COST Action IC1104 and grant KU 2430/3-1 and WA 1666/9-1 – Integer Linear Programming Models for Subspace Codes and Finite Geometry from the German Research Foundation.

dimension case $K = \{k\}$, respectively. Note that in the latter case $d_S(U,W) = 2 \cdot d_I(U,W) \in 2 \cdot \mathbb{N}$ is an even number.

In general, the exact determination of $A_q^S(n,d;K)$ is a hard problem, both on the theoretic and the algorithmic side. Therefore, lower and upper bounds on $A_q^S(n,d;K)$ have been intensively studied in the last years, see e.g. [7]. Since the underlying discrete structures arose under different names in different fields of discrete mathematics, it is even more difficult to get an overview of the state of the art. For example, geometers are interested in so-called partial $(k-1)$-spreads of $\mathrm{PG}(n-1,q)$. Following the track of partial spreads, one can end up with orthogonal arrays or $(s,r,\mu)$-nets. Furthermore, $q$-analogs of Steiner systems provide optimal constant dimension codes. For some sets of parameters constant dimension codes are in one-to-one correspondence with so-called vector space partitions.

The aim of this report is to describe the underlying theoretical base of an on-line database[1] maintained by the authors that tries to collect up-to-date information on the best lower and upper bounds for subspace codes. Whenever the exact value $A_q^S(n,d;K)$ could be determined, we ask for a complete classification of all optimal codes up to isomorphism. Since the overall task is rather comprehensive, we start by focusing on the special cases of constant dimension codes, $A_q^S(n,d;k)$, and (unrestricted) subspace codes, $A_q^S(n,d)$, using the subspace distance as metric. For a more comprehensive survey on network coding we refer the interested reader e.g. to [2]. For algorithmic aspects we refer the interested reader e.g. to [16].

The remaining part of this report is structured as follows. In Section 2 we outline how to use the tables. The currently implemented constructions and upper bounds are described in Section 3 and Section 4, respectively. The still rather unsteady Application programming interface (API) is the topic of Section 5. Finally we draw a conclusion in Section 6 and list some explicit tables on upper and lower bounds in an appendix.

## 2. How to use the tables

On the website the two special cases $A_q^S(n,d;k)$ and $A_q^S(n,d)$ can be accessed via the menu items `CDC` (constant dimension code) and `MDC` (mixed dimension code), see Figure 1. Selecting the item `Table` yields the rough data that we will outline in this section. Selecting the item `Constraints` yields information abound the so far implemented general-purpose lower bounds, see Section 3, and upper bounds, see Section 4.

### 2.1. Constant dimension codes – CDC.

For a constant dimension code the dimension $n$ of the ambient space (first *selection row*) and the field size $q$ (second *selection row*) can be chosen. The current limits are $4 \leq n \leq 19$ and $2 \leq q \leq 9$. For each chosen pair of those parameters a table with the information on lower and upper bounds on constant dimension codes over $\mathbb{F}_q^n$ is displayed.

The rows of those tables are labeled by the minimum distance $d = d_S(\star)$ and the columns are labeled by the dimension $k$ of the codewords. In the third *selection row* several *views* can be picked. The first three options, `short`, `normal`, and `large`, specify the subset of possible values for the parameters $d$ and $k$. In the most extensive view `large`, $k$ can take all integers between 0 and $n$. For $d$ the integers between 1 and $n$ are considered. As

- $A_q^S(n,d;0) = 1$ for all $1 \leq d \leq n$;
- $A_q^S(n,d;k) = A_q^S(n,d;n-k)$;
- $A_q^S(n,2d'+1;k) = A_q^S(n,2d'+2;k)$ for all $d' \in \mathbb{N}_{>0}$;

one may assume $1 \leq k \leq \lfloor n/2 \rfloor$, $2 \leq d \leq n$, and $d \in 2\mathbb{N}$. These assumptions are implemented in the view `normal`. However, some exact values $of A_q^S(n,d;k)$ are rather easy to determine

- $A_q^S(n,2;k) = \begin{bmatrix} n \\ k \end{bmatrix}_q$, since any two different $k$-dimensional subspaces of $\mathbb{F}_q^n$ have a subspace distance of at least 2;

[1] http://subspacecodes.uni-bayreuth.de


subspacecodes.uni-bayreuth.de
subspacecodes.uni-bayreuth.de/table/2/10/short/
SCT
CDC
MDC
Literature
About
Contribute
n=4 n=5 n=6 n=7 n=8 n=9 n=10 n=11 n=12 n=13 n=14 n=15 n=16 n=17 n=18 n=19
q=2 q=3 q=4 q=5 q=7 q=8 q=9
short normal large - gap density - amount mrd bound amount pending dots amount lifted mrd
Table for $A_2(10, d; k)$
d\k 2 3 4 5
4 341 23870 - 24698 301213 - 423181 1167355 - 1678413
6 145 4173 - 4978 32843 - 38214
8 65 1025 - 1089
10 33

FIGURE 1. Tables of constant dimension codes

- if $d > 2k$, then we can have at most one codeword, i.e., $A_q^S(n, d; k) = 1$.

Thus, we may assume $2 \leq k \leq \lfloor n/2 \rfloor$, $4 \leq d \leq 2k$, and $d \in 2\mathbb{N}$. These assumptions are implemented in the view `short`. The standard selection is given by $n = 4$, $q = 2$ and view `short`.

Given one of these three views, a table entry may consist of

- a range $l$–$u$: An example is given by the parameters $q = 2$, $n = 7$, $d = 4$, $k = 3$, where $l = 329$ and $u = 381$. The meaning is that for the corresponding maximum cardinality of a constant dimension code only the lower bound $l$ and the upper bound $u$ is known, i.e., $329 \leq A_2^S(7, 4; 3) \leq 381$ in the example.
- a **bold** number $m$: An example is given by the parameters $q = 2$, $n = 10$, $d = 8$, $k = 4$, where $m = 65$. The meaning is that the corresponding maximum cardinality of a constant dimension code is exactly determined, i.e., $A_2^S(10, 8; 4) = 65$ in the example.
- a **bold** number $m$ with an asterisk and a number $l$ in brackets: An example is given by the parameters $q = 2$, $n = 6$, $d = 4$, $k = 3$, where $m = 77$ and $l = 5$. The meaning is that the corresponding maximum cardinality of a constant dimension code is exactly determined and all optimal codes have been classified up to isomorphism, i.e., $A_2^S(6, 4; 3) = 77$ and there are exactly 5 isomorphism types in the example [14]. Another example is given for the parameters $q = 2$, $n = 6$, $d = 4$, and $k = 2$, where there are exactly $131,044$ isomorphism types of constant dimension codes attaining cardinality $A_2^S(6, 4; 2) = 21$ [18].

Each nontrivial table entry is clickable and then yields further information on several lower and upper bounds, see Section 3 and Section 4 for the details.

In some cases, e.g., for the parameters $q = 2$, $n = 6$, $d = 4$, and $k = 3$, the corresponding codes are also available for download using the button called "file". The format of these codes is GAP[2].

Besides the views `short`, `normal`, and `large` for the selection of ranges for the parameters $d$ and $k$, there are some additional views. The views `relative gap` and `ratio of bounds` condense the current lack of knowledge on the exact value of $A_q^S(n, d; k)$ to a single number. For the view `relative gap` this number is given by the formula

$$\frac{\text{upper bound} - \text{lower bound}}{\text{lower bound}},$$

i.e., we obtain a non-negative real number. While principally any number in $\mathbb{R}_{\geq 0}$ can be obtained, the largest relative gap in our database is currently given by about $0.728$ for the parameters $q = 2, n = 19, d = 4, k = 9$. A gap of $0.0$ corresponds to the determination of the exact value $A_q^S(n, d; k)$. The mentioned formula is also displayed on the webpage, when you move your mouse over the word `relative gap`. For the view `ratio of bounds` the corresponding number is given by the formula

$$\frac{\text{lower bound}}{\text{upper bound}},$$

which may take any real number in $(0, 1]$. The smallest ratio of bounds in our database is given by about $0.578$ for the same parameters as above. Clearly, the largest relative gap yields the smallest ratio of bounds and vice versa as the function $x \mapsto \frac{1}{x} - 1$ is strictly decreasing in $(0, 1]$. A ratio of bounds of $1.0$ corresponds to the determination of the exact value $A_q^S(n, d; k)$. The mouse-over effect is also implemented in that case.

Another type of views arose from some of the various constructions described in Section 3. They are all labeled as `amount pending dots` and `amount lifted mrd` and condense the *strength* of a certain construction to a single number in $\mathbb{R}_{\geq 1}$. This number is always given as the quotient between the currently best known lower bound and the value obtained by the respective construction. Here, a value of one means that the currently best known code can be obtained by the respective construction. A value larger than $1$ measures how much better a more tailored construction is for this specific set of parameters compared to the respective general construction method. We remark that `amount pending dots` is still experimental and in some cases there may still be better codes obtained from the underlying very general construction technique, which has quite some degrees of freedom. With respect to upper bounds the additional view `amount mrd bound` is introduced. Here the displayed single number is given by the currently best known lower bound divided by the so-called MRD bound, see Subsection 4.2.

2.2. **Mixed dimension codes – `MDC`.** For a subspace code with mixed dimensions the field size $q$ (*selection row* number one) can be chosen. The current limits are given by $2 \leq q \leq 9$. For each chosen parameter a table with the information on lower and upper bounds on mixed dimension codes over $\mathbb{F}_q^n$ is displayed, see Figure 2.

The rows of those tables are labeled by the distance $d = d_S(\star)$ and the columns are label by the dimension $n$ of the ambient space $\mathbb{F}_q^n$. In the second *selection row* several *views* can be picked. The view `normal`, c.f. Subsection 2.1, already incorporates the restriction to $1 \leq d \leq n \leq 19$. The views `relative gap` and `ratio of bounds` condense the current lack of knowledge on the exact value of $A_q^S(n, d)$ to a single number. For the view `relative gap` this number is given by the formula

$$\frac{\text{upper bound} - \text{lower bound}}{\text{lower bound}},$$

i.e. we obtain a non-negative real number. While principally any number in $\mathbb{R}_{\geq 0}$ can be obtained, the largest relative gap in our database is currently given by about $2.899$ for the parameters $q = 2, n =$

[2] `http://www.gap-system.org`




FIGURE 2. Tables of subspace codes

$19, d = 4$. A relative gap of $0.0$ corresponds to the determination of the exact value $A_q^S(n, d)$. The mentioned formula is also displayed on the webpage, when you move your mouse over the word `relative gap`. For the view `ratio of bounds` the corresponding number is given by the formula

$$\frac{\text{lower bound}}{\text{upper bound}},$$

which may take any real number in $(0, 1]$. The smallest ratio of bounds in our database is given by about $0.256$ for the same parameters as above. Clearly the largest relative gap yields the smallest ratio of bounds and vice versa as the function $x \mapsto \frac{1}{x} - 1$ is strictly decreasing in $(0, 1]$. A ratio of bounds of $1.0$ corresponds to the determination of the exact value $A_q^S(n, d)$. The mouse-over effect is also implemented in that case.

## 3. IMPLEMENTED CONSTRUCTIONS – LOWER BOUNDS

3.1. **Lifted MRD codes.** For matrices $A, B \in \mathbb{F}_q^{m\times n}$ the rank distance is defined via $d_R(A, B) := \mathrm{rk}(A - B)$. It is indeed a metric, as observed in [12].

**Theorem 3.1.** *(see* [12]*) Let $m, n \geq d$ be positive integers, $q$ a prime power, and $\mathcal{C} \subseteq \mathbb{F}_q^{m\times n}$ be a rank-metric code with minimum rank distance $d$. Then, $|\mathcal{C}| \leq q^{\max(n,m)\cdot(\min(n,m)-d+1)}$.*

Codes attaining this upper bound are called maximum rank distance (MRD) codes. They exist for all (suitable) choices of parameters. If $m < d$ or $n < d$, then only $|\mathcal{C}| = 1$ is possible, which may be summarized to the single upper bound $|\mathcal{C}| \leq \left\lceil q^{\max(n,m)\cdot(\min(n,m)-d+1)} \right\rceil$. Using an $m \times m$ identity matrix as a prefix one obtains the so-called lifted MRD codes.

**Theorem 3.2.** *(see* [21]*) For positive integers $k, d, n$ with $k \leq n$, $d \leq 2\min(k, n-k)$, and $d \equiv 0 \pmod 2$, the size of a lifted MRD code in $G_q(n,k)$ with subspace distance $d$ is given by*

$$M(q,k,n,d) := q^{\max(k,n-k)\cdot(\min(k,n-k)-d/2+1)}.$$

*If $d > 2\min(k, n-k)$, then we have $M(q,k,n,d) = 1$.*

3.2. **Echelon-Ferrers or multilevel construction.** In [8] a generalization, the so-called multi-level construction, based on lifted MRD codes was presented. Let $1 \leq k \leq n$ be integers and $v \in \mathbb{F}_2^n$ a binary vector of weight $k$. By $\mathrm{EF}_q(v)$ we denote the set of all $k \times n$ matrices over $\mathbb{F}_q$ that are in row-reduced echelon form, i.e. the Gaussian algorithm had been applied, and the pivot columns coincide with the positions where $v$ has a 1-entry.

**Theorem 3.3.** *(see* [8]*) For integers $k, n, \delta$ with $1 \leq k \leq n$ and $1 \leq \delta \leq \min(k, n-k)$, let $\mathcal{B}$ be a binary constant weight code of length $n$, weight $k$, and minimum Hamming distance $2\delta$.*

For each $b \in \mathcal{B}$ let $\mathcal{C}_b$ be a code in $\mathrm{EF}(b)$ with minimum rank distance at least $\delta$. Then, $\cup_{b\in\mathcal{B}} \mathcal{C}_b$ is a constant dimension code of dimension $k$ having a subspace distance of at least $2\delta$.

**Theorem 3.4.** *(see* [8]*) Let $\mathcal{F}$ be the Ferrers diagram of $\mathrm{EF}_q(v)$ and $\mathcal{C} \subseteq \mathrm{EF}_q(v)$ be a subspace code having a subspace distance of at least $2\delta$, then*

$$|\mathcal{C}| \leq q^{\min_{0\leq i\leq \delta-1} \nu_i},$$

*where $\nu_i$ is the number of dots in $\mathcal{F}$, which are neither contained in the first $i$ rows nor contained in the rightmost $\delta - 1 - i$ columns.*

The authors of [8] conjecture that Theorem 3.4 is tight for all parameters $q$, $\mathcal{F}$, and $\delta$, which is still unrebutted.

Taking binary vectors with $k$ consecutive ones we are in the classical MRD case. So, taking binary vectors $v_i$, where the ones are located in positions $(i-1)k+1$ to $ik$ for all $1 \leq i \leq \lfloor n/k \rfloor$, clearly gives a binary constant weight code of length $n$, weight $k$, and minimum Hamming distance $2k$.

**Observation 3.5.** *(see e.g.* [17]*) For positive integers $k$, $n$ with $n > 2k$ and $n \not\equiv 0 \pmod k$, there exists a constant dimension code in $G_q(n,k)$ with subspace distance $2k$ having cardinality*

$$1 + \sum_{i=1}^{\lfloor n/k \rfloor - 1} q^{n-ik} = 1 + q^{k+(n \bmod k)} \cdot \frac{q^{n-k-(n \bmod k)} - 1}{q^k - 1} = \frac{q^n - q^{k+(n \bmod k)} + q^k - 1}{q^k - 1}.$$

We remark that a more general construction, among similar lines and including explicit formulas for the respective cardinalities, has been presented in [22].

We remark that the general Echelon-Ferrers or multilevel construction contains the mentioned observation as a very easy special case. However, our knowledge on the size of an MRD code over $\mathrm{EF}(v)$ is still very limited. As mentioned, there is an explicit conjecture, which so far is neither proven nor disproven. The construction has even been fine-tuned to the so-called pending dots [9, 19] and the so-called pending blocks [20] constructions.

3.3. **Nonconstructive bounds.**

**Theorem 3.6.** *(Sphere covering, see* [15]*)*

$$A_q^S(n,d;k) \geq \begin{bmatrix} n \\ k \end{bmatrix}_q \Big/ \left( \sum_{i=0}^{(d/2-1)+1} \begin{bmatrix} k \\ i \end{bmatrix}_q \cdot \begin{bmatrix} n-k \\ i \end{bmatrix}_q \cdot q^{i^2} \right)$$

This lower bound is implemented as `sphere_covering`.

### 3.4. **Explicit, nonrecurring constructions.**

**Theorem 3.7.** *(Graham, Sloane, see* [23]*)*

$$A_q^S(n,d;k) \geq \frac{(q-1)\begin{bmatrix} n \\ k \end{bmatrix}_q}{(q^n-1)q^{n(d/2-2)}}$$

This lower bound is implemented as `graham_sloane`.

**Theorem 3.8.** *(Linearized polynomials, see* [15]*)*

$$A_q^S(n,d;k) \geq q^{(n-k)(k-d/2+1)}$$

This lower bound is implemented as `lin_poly`.

**Theorem 3.9.** *(Partial spreads, see* [11]*) If $d = 2k$ then:*

$$A_q^S(n,d;k) \geq \frac{q^n - q^k(q^{(n \bmod k)}-1)-1}{q^k-1}$$

This lower bound is implemented as `partial_spread_3`.

## 4. Implemented upper bounds

Assuming $0 \leq k \leq n$ we always have $A_q^S(n,d;k) \geq 1$. Since we can take no more than all subspaces of a given dimension, we obtain the trivial upper bound $A_q^S(n,d;k) \leq \begin{bmatrix} n \\ k \end{bmatrix}_q$ which is implemented as `all_subs`.

### 4.1. ***Classical*** **coding theory bounds.**

**Theorem 4.1.** *(Singleton bound, see* [15]*)*

$$A_q^S(n,d;k) \leq \begin{bmatrix} n-d/2+1 \\ k-d/2+1 \end{bmatrix}_q$$

This upper bound is implemented as `singleton`.

**Theorem 4.2.** *(Sphere packing bound, see* [15]*)*

$$A_q^S(n,d;k) \leq \begin{bmatrix} n \\ k \end{bmatrix}_q \Bigg/ \left( \sum_{i=0}^{(d/2-1)/2+1} \begin{bmatrix} k \\ i \end{bmatrix}_q \cdot \begin{bmatrix} n-k \\ i \end{bmatrix}_q \cdot q^{i^2} \right)$$

This upper bound is implemented as `sphere_packing`.

**Theorem 4.3.** *(Johnson bounds, see* [11]*)*

$$A_q^S(n,d;k) \leq \left\lfloor \frac{(q^n-1)\cdot A_q^S(n-1,d;k-1)}{q^k-1} \right\rfloor$$

*and*

$$A_q^S(n,d;k) \leq \left\lfloor \frac{(q^n-1)\cdot A_q^S(n-1,d;k)}{q^{n-k}-1} \right\rfloor$$

These upper bounds are implemented as `johnson_1` and `johnson_2`.

**Theorem 4.4.** *(Anticode bounds, see* [11]*)*

$$A_q^S(n,d;k) \leq \left\lfloor \begin{bmatrix} n \\ k \end{bmatrix}_q \Bigg/ \begin{bmatrix} n-k+d/2-1 \\ d/2-1 \end{bmatrix}_q \right\rfloor$$

This upper bound is implemented as `anticode`.

4.2. **MRD bound.** Since the size of the lifted MRD code, see Theorem 3.1, is quite competitive it is interesting to compare the best known constructions with this very general explicit construction. Even more, lifted MRD codes are the basis for more involved constructions, see Subsection 3.2. From this point of view it is very interesting that an upper bound for the cardinality of constant dimension codes containing the lifted MRD code (of shape $k \times (n-k)$ and rank distance $d/2$) can be stated that is tighter than the best known general upper bounds:

**Theorem 4.5.** *(see* [9, Theorem 10 and 11]*) Let $\mathcal{C} \subseteq G_n(k,q)$, where $n \geq 2k$, with minimum subspace distance $d$ that contains the lifted MRD code.*

- *If $d = 2(k-1)$ and $k \geq 3$, then $|\mathcal{C}| \leq q^{2(n-k)} + A_q(n-k, 2(k-2); k-1)$;*
- *if $d = k$, where $k$ is even, then $|\mathcal{C}| \leq q^{(n-k)(k/2+1)} + \begin{bmatrix} n-k \\ k/2 \end{bmatrix}_q \frac{q^n - q^{n-k}}{q^k - q^{k/2}} + A_q(n-k, k; k)$.*

4.3. **Bounds for spreads.**

**Theorem 4.6.** *(*[1]*; see also* [4, p. 29]*, Result 2.1 in* [3]*) $\mathbb{F}_q^n$ contains a $k$-spread if and only if $k$ divides $n$, where we assume $1 \leq k \leq n$ and $k, n \in \mathbb{N}$.*

The corresponding exact value is implemented as upper bound `spread`.

**Theorem 4.7.** *(*[3]*; see also* [13] *for the special case $q = 2$) For positive integers $1 \leq k \leq n$ be positive integers with $n \equiv 1 \pmod{k}$ we have $A_q^S(n, 2k; k) = \frac{q^n - q}{q^k - 1} - q + 1 = q \cdot \frac{q^{n-1} - 1}{q^k - 1} - q + 1 = \frac{q^n - q^{k+1} + q^k - 1}{q^k - 1}$.*

The corresponding exact value is implemented as upper bound `partial_spread_2`.

**Theorem 4.8.** *(Corollary 8 in* [5]*) If $n = k(t+1) + r$ with $0 < r < k$, then*

$$A_q^S(n, 2k; k) \leq \sum_{i=0}^{t} q^{ik+r} - \lfloor \theta \rfloor - 1 = q^r \cdot \frac{q^{k(t+1)} - 1}{q^k - 1} - \lfloor \theta \rfloor - 1,$$

*where $2\theta = \sqrt{1 + 4q^k(q^k - q^r)} - (2q^k - 2q^r + 1)$.*

We remark that this theorem is also restated as Theorem 13 in [7] and as Theorem 44 in [10] with the small typo of not rounding down $\theta$ ($\Omega$ in their notation). The corresponding upper bound is implemented as `partial_spread_4`.

**Theorem 4.9.** *(see* [6]*) For each integer $m \geq 2$ we have*

(a) $A_2(3m, 6; 3) = \frac{2^{3m} - 1}{7}$;
(b) $A_2(3m+1, 6; 3) = \frac{2^{3m+1} - 9}{7}$;
(c) $A_2(3m+2, 6; 3) = \frac{2^{3m+2} - 18}{7}$.

The corresponding upper bound is implemented as `partial_spread_1`.

**Theorem 4.10.** *(Theorem 4.3 in* [17]*) For each pair of integers $t \geq 1$ and $k \geq 4$ we have $A_2^S(k(t+1) + 2, 2k; k) = \frac{2^{k(t+1)+2} - 3 \cdot 2^k - 1}{2^k - 1}$.*

The corresponding upper bound is implemented as `partial_spread_kurz_q2`.

**Lemma 4.11.** *(Lemma 4.6 in* [17]*) For integers $t \geq 1$ and $k \geq 4$ we have $A_3^S(k(t+1) + 2, 2k; k) \leq \frac{3^{k(t+1)+2} - 3^2}{3^k - 1} - \frac{3^2 + 1}{2}$.*

The corresponding upper bound is implemented as `partial_spread_kurz_q3`.

## 5. APPLICATION PROGRAMMING INTERFACE

There is also an API available to access most data of the database. It is inspired by the REST (representational state transfer) style and only GET querys are supported. In order to access the data for the constant dimension case with parameters $q, n, d$ and $k$, you query the URL

http://subspacecodes.uni-bayreuth.de/api/$q$/$n$/$d$/$k$/

Similarly in the mixed dimension case, the URL is

http://subspacecodes.uni-bayreuth.de/api/$q$/$n$/$d$/

The result is a JSON file which contains a subset of the following attributes:

- request = contains your specified $q, n, d$ and $k$
- $\{$lower,upper$\}$_bound = lower or upper bound for the value $A_q(n, d; k)$
- comments = commentaries to this entry
- nondeduced = if the parameters are no parameters that are also viewable in the "short" mode, then they are trivial or computed using other parameters. nondeduced lists these other parameters.
- $\{$lower,equal,upper$\}$_bound_constraints = list of tuples which contain name, parameter and value of the applied constraints
- classified = boolean that is true if $A_q(n, d; k)$ is classified up to isomorphism
- known codes = list of tuples of size, details, file (to enable automatic downloads) and nrisotypes (the number of isomorphism types of this entry)
- liftedmrdsizebound = the bound for codes that contains the lifted MRD code as described in 4.2

In order to download the codes, you have to use the attribute file above and the URL

http://subspacecodes.uni-bayreuth.de/codes/*file*

We want to remark that the API (as well as the whole homepage) is still in an early evolutionary phase and therefore changes are likely to occur. As an example, the URL

`http://subspacecodes.uni-bayreuth.de/api/2/6/4/3/`

yields the output:

{"upper_bound_constraints": [{"parameter": "", "name": "all_subs", "value": 1395},
{"parameter": "", "name": "anticode", "value": 93}, {"parameter": "2", "name": "ilp_2", "value": 93},
{"parameter": "4", "name": "ilp_3", "value": 93}, {"parameter": "", "name": "singleton", "value": 155},
{"parameter": "", "name": "sphere_packing", "value": 1395}, {"parameter": "1", "name": "ilp_1", "value": 81},
{"parameter": "5", "name": "ilp_4", "value": 81}, {"parameter": "", "name": "johnson_1", "value": 81},
{"parameter": "", "name": "johnson_2", "value": 81}],
"known_codes": [{"nrisotypes": 5, "details": "", "file": "code_2_6_4_3_optimal_size_77.zip", "size": 77}], "upper_bound": 77,
"classified": true, "lower_bound": 77, "lower_bound_constraints": [{"parameter": "", "name": "HonoldKiermaierKurz_n6_d4_k3",
"value": 77}, {"parameter": "", "name": "construction_1", "value": 71}, {"parameter": "", "name": "sphere_covering",
"value": 15}, {"parameter": "", "name": "trivial_1", "value": 0}, {"parameter": "3", ], "request": [2, 6, 4, 3], "liftedmrdsize-
bound": 71, "comments": "", "equal_bound_constraints": []}

## 6. CONCLUSION

The collection of the known results on lower and upper bounds for subspace codes is an ongoing project. So far we have merely implemented the tip of the iceberg of the available knowledge. We still hope that the emerging on-line data base and the accompanying user's guide is already valuable for researcher in the field at this early stage. If you want to support us in this task, please let us know any known constructions, bounds or papers that we have missed so far via daniel.heinlein@uni-bayreuth.de or the *contribute*-button in the upper right corner of the webpage `subspacecodes.uni-bayreuth.de`.

Tracing back results to their original source is a task on its own. We want to work on that issue more intensively in the future. If you observe possible enhancements in that direction, please let us know. Critique, suggestions for improvements and feature requests are also highly welcome.

## REFERENCES


[1] J. André, *Über nicht-desarguessche Ebenen mit transitiver Translationsgruppe*, Mathematische Zeitschrift **60** (1954), no. 1, 156–186.

[2] R. Bassoli, H. Marques, J. Rodriguez, K.W. Shum, and R. Tafazolli, *Network coding theory: A survey*, Communications Surveys & Tutorials, IEEE **15** (2013), no. 4, 1950–1978.

[3] A. Beutelspacher, *Partial spreads in finite projective spaces and partial designs*, Mathematische Zeitschrift **145** (1975), no. 3, 211–229.

[4] P. Dembowski, *Finite Geometries: Reprint of the 1968 edition*, Springer Science & Business Media, 2012.

[5] D.A. Drake and J.W. Freeman, *Partial $t$-spreads and group constructible $(s, r, \mu)$-nets*, Journal of Geometry **13** (1979), no. 2, 210–216.

[6] S. El-Zanati, H. Jordon, G. Seelinger, P. Sissokho, and L. Spence, *The maximum size of a partial 3-spread in a finite vector space over $GF(2)$*, Designs, Codes and Cryptography **54** (2010), no. 2, 101–107.

[7] T. Etzion, *Problems on q-analogs in coding theory*, arXiv preprint: 1305.6126, 37 pages (2013).

[8] T. Etzion and N. Silberstein, *Error-correcting codes in projective spaces via rank-metric codes and Ferrers diagrams*, IEEE Transactions on Information Theory **55** (2009), no. 7, 2909–2919.

[9] ———, *Codes and designs related to lifted MRD codes*, IEEE Transactions on Information Theory **59** (2013), no. 2, 1004–1017.

[10] T. Etzion and L. Storme, *Galois geometries and coding theory*, Designs, Codes and Cryptography (to appear), 49p.

[11] T. Etzion and A. Vardy, *Error-correcting codes in projective space*, IEEE Transactions on Information Theory **57** (2011), no. 2, 1165–1173.

[12] E.M. Gabidulin, *Theory of codes with maximum rank distance*, Problemy Peredachi Informatsii **21** (1985), no. 1, 3–16.

[13] S.J. Hong and A.M. Patel, *A general class of maximal codes for computer applications*, IEEE Transactions on Computers **100** (1972), no. 12, 1322–1331.

[14] T. Honold, M. Kiermaier, and S. Kurz, *Optimal binary subspace codes of length 6, constant dimension 3 and minimum distance 4*, Contemp. Math. **632** (2015), 157–176.

[15] R. Koetter and F.R. Kschischang, *Coding for errors and erasures in random network coding*, IEEE Transactions on Information Theory **54** (2008), no. 8, 3579–3591.

[16] A. Kohnert and S. Kurz, *Construction of large constant dimension codes with a prescribed minimum distance*, Mathematical methods in computer science, Springer, 2008, pp. 31–42.

[17] S. Kurz, *Improved upper bounds for partial spreads*, arXiv preprint: 1512.04297, 8 pages (2015), 8.

[18] Zlatka T. Mateva and Svetlana T. Topalova, *Line spreads of $\mathrm{PG}(5, 2)$*, J. Combin. Des. **17** (2009), no. 1, 90–102. MR 2475427 (2010e:51008)

[19] N. Silberstein and T. Etzion, *Codes and designs related to lifted MRD codes*, Information Theory Proceedings (ISIT), 2011 IEEE International Symposium on, IEEE, 2011, pp. 2288–2292.

[20] N. Silberstein and A.-L. Trautmann, *Subspace codes based on graph matchings, ferrers diagrams and pending blocks*, arXiv preprint: 1404.6723, pages 25 (2014).

[21] D. Silva, F.R. Kschischang, and R. Koetter, *A rank-metric approach to error control in random network coding*, IEEE Transactions on Information Theory **54** (2008), no. 9, 3951–3967.

[22] V. Skachek, *Recursive code construction for random networks*, IEEE transactions on Information Theory **56** (2010), no. 3, 1378–1382.

[23] S.-T. Xia, *A graham-sloane type construction of constant dimension codes*, Network Coding, Theory and Applications, 2008. NetCod 2008. Fourth Workshop on, IEEE, 2008, pp. 1–5.

## APPENDIX A. TABLES FOR BINARY CONSTANT DIMENSION CODES

| $n=6$ | 2 | 3 |
|---|---|---|
| 4 | 21 * (131044) | 77 * (5) |
| 6 | | 9 |

| $n=7$ | 2 | 3 |
|---|---|---|
| 4 | 41 | 329–381 |
| 6 | | 17 |

| $n=8$ | 2 | 3 | 4 |
|---|---|---|---|
| 4 | 85 | 1326–1493 | 4801–6477 |
| 6 | | 34 | 257–289 |
| 8 | | | 17 |

| $n=9$ | 2 | 3 | 4 |
|---|---|---|---|
| 4 | 169 | 5986–6205 | 37265–50861 |
| 6 | | 73 | 1033–1158 |
| 8 | | | 33 |

| $n=10$ | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| 4 | 341 | 23870–24698 | 301213–423181 | 1167355–1678413 |
| 6 | | 145 | 4173–4978 | 32843–38214 |
| 8 | | | 65 | 1025–1089 |
| 10 | | | | 33 |

| $n=11$ | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| 4 | 681 | 97526–99718 | 2134417–3370453 | 16814481–27943597 |
| 6 | | 290 | 16641–19787 | 262780–328708 |
| 8 | | | 129–133 | 4097–4292 |
| 10 | | | | 65 |

| $n=12$ | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|
| 4 | 1365 | 385515–398385 | 19664917–27223014 | 269602811–445225968 | 1074043037–1816333805 |
| 6 | | 585 | 66569–79170 | 2098185–2613798 | 16777289–21366020 |
| 8 | | | 273 | 16385–17568 | 262145–278980 |
| 10 | | | | 129 | 4097–4225 |
| 12 | | | | | 65 |

| $n=13$ | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|
| 4 | 2729 | 1597245 | 157319497–217544769 | 4311781777–7193022828 | 34376552849–57886442918 |
| 6 | | 1169 | 266891–319449 | 16810059–20918757 | 268439629–339835228 |
| 8 | | | 545 | 65569–72133 | 2097153–2284118 |
| 10 | | | | 257–260 | 16385–16772 |
| 12 | | | | | 129 |

## APPENDIX B. TABLES FOR TERNARY CONSTANT DIMENSION CODES

| $n=6$ | 2 | 3 |
|---|---|---|
| 4 | 91 | 754–784 |
| 6 | | 28 |

| $n=7$ | 2 | 3 |
|---|---|---|
| 4 | 271 | 6977–7651 |
| 6 | | 82 |

| $n=8$ | 2 | 3 | 4 |
|---|---|---|---|
| 4 | 820 | 60259–68375 | 543142–627382 |
| 6 | | 244–248 | 6562–6724 |
| 8 | | | 82 |

| $n=9$ | 2 | 3 | 4 |
|---|---|---|---|
| 4 | 2458 | 549667–620740 | 14585908–16821959 |
| 6 | | 757 | 59059–61014 |
| 8 | | | 244 |

| $n=10$ | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| 4 | 7381 | 5086234–5582307 | 394218370–458168194 | 3539131096–4104557996 |
| 6 | | 2269 | 531689–558741 | 14349658–14887416 |
| 8 | | | 730–733 | 59050–59536 |
| 10 | | | | 244 |

| $n=11$ | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| 4 | 22141 | 45776125–50289024 | 10474939111–12361041947 | 282444122389–335382904522 |
| 6 | | 6805–6809 | 4789531–5024303 | 387441117–409003029 |
| 8 | | | 2188–2202 | 531442–536562 |
| 10 | | | | 730 |

## APPENDIX C. TABLES FOR QUATERNARY CONSTANT DIMENSION CODES

| $n=6$ | 2 | 3 |
|---|---|---|
| 4 | 273 | 4137–4225 |
| 6 | | 65 |

| $n=7$ | 2 | 3 |
|---|---|---|
| 4 | 1089 | 66828–70993 |
| 6 | | 257 |

| $n=8$ | 2 | 3 | 4 |
|---|---|---|---|
| 4 | 4369 | 1054373–1132819 | 16874321–18245201 |
| 6 | | 1025–1033 | 65537–66049 |
| 8 | | | 257 |

| $n=9$ | 2 | 3 | 4 |
|---|---|---|---|
| 4 | 17473 | 16945153–18179409 | 1078557605–1164551259 |
| 6 | | 4161 | 1048593–1061936 |
| 8 | | | 1025 |

| $n=10$ | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| 4 | 69905 | 273727489–290821444 | 69040760145–74754799185 | 1104214839637–1193665040475 |
| 6 | | 16641 | 16778246–17110276 | 1073745957–1088484400 |
| 8 | | | 4097–4104 | 1048577–1050625 |
| 10 | | | | 1025 |

| $n=11$ | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| 4 | 279617 | 4379639873–4654011924 | 4399125068709–4783502960915 | 281476055268261–306494895880785 |
| 6 | | 66561–66569 | 268500993–273715279 | 68719743536–70152181776 |
| 8 | | | 16385–16418 | 16777217–16826412 |
| 10 | | | | 4097 |

## APPENDIX D. TABLE FOR (UNRESTRICTED) BINARY SUBSPACE CODES

| $q=2$ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 2 | 5 | 16 | 67 | 374 | 2825 | 29212 | 417199 | 8283458 | 229755605 |
| 2 | | 3 | 8 | 37 | 187 | 1521 | 14606 | 222379 | 4141729 | 121919127 |
| 3 | | | 2 | 5 | 18 | 104–118 | 593–776 | 4907–9268 | 70748–107419 | 1242813–2532929 |
| 4 | | | | 5 | 9 | 77 | 330–463 | 4803–9268 | 37267–107419 | 1167613–2532929 |
| 5 | | | | | 2 | 9 | 34 | 261–359 | 1990–2462 | 32971–48402 |
| 6 | | | | | | 9 | 17 | 257–357 | 1033–2462 | 32843–48402 |
| 7 | | | | | | | 2 | 17 | 62–66 | 1025–1219 |
| 8 | | | | | | | | 17 | 33 | 1025–1219 |
| 9 | | | | | | | | | 2 | 33 |
| 10 | | | | | | | | | | 33 |

## APPENDIX E. TABLE FOR (UNRESTRICTED) TERNARY SUBSPACE CODES

| $q=3$ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
|---|---|---|---|---|---|---|---|---|---|
| 1 | 2 | 6 | 28 | 212 | 2664 | 56632 | 2052656 | 127902864 | 13721229088 |
| 2 | | 4 | 14 | 132 | 1332 | 34608 | 1026328 | 77705744 | 6860614544 |
| 3 | | | 2 | 10 | 53–56 | 764–966 | 13247–15802 | 544431–762424 | 29009271–34890316 |
| 4 | | | | 10 | 28 | 754–966 | 6978–15802 | 543144–762424 | 14585910–34890314 |
| 5 | | | | | 2 | 28 | 160–164 | 6568–7222 | 117620–123544 |
| 6 | | | | | | 28 | 82 | 6562–7220 | 59060–123542 |
| 7 | | | | | | | 2 | 82 | 484–488 |
| 8 | | | | | | | | 82 | 244 |
| 9 | | | | | | | | | 2 |

## APPENDIX F. TABLE FOR (UNRESTRICTED) QUATERNARY SUBSPACE CODES

| $q=4$ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|---|---|---|---|---|---|---|---|---|
| 1 | 2 | 7 | 44 | 529 | 12278 | 565723 | 51409856 | 9371059621 |
| 2 | | 5 | 22 | 359 | 6139 | 379535 | 25704928 | 6269331761 |
| 3 | | | 2 | 17 | 127–130 | 4154–4771 | 131317–144060 | 16881731–20519579 |
| 4 | | | | 17 | 65 | 4137–4771 | 66829–144060 | 16874323–20519577 |
| 5 | | | | | 2 | 65 | 510–514 | 65545–68117 |
| 6 | | | | | | 65 | 257 | 65537–68115 |
| 7 | | | | | | | 2 | 257 |
| 8 | | | | | | | | 257 |

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BAYREUTH, 95440 BAYREUTH, GERMANY
*E-mail address*: firstname.lastname@uni-bayreuth.de